Corrected Table 5.4: Clinical Institute Withdrawal Assessment (CIWA) CARPA STM 8th edition page 281

 Nausea and vomiting No nausea and no vomiting Mild nausea and no vomiting Mild nausea and no vomiting Intermittent nausea, with dry retching Constant nausea, frequent dry retching or vomiting 		 2. Tremor 0. No tremor 1. Not visible, but can be felt fingertip to fingertip 2. 3. 4. Moderate 5. 6. 7. Severe, even with arms not extended 		 3. Sweating 0. No sweat visible 1. Barely perceptible sweating, palms moist 2. 3. 4. Beads of sweat obvious on forehead 5. 6. 7. Drenching sweats
 4. Anxiety 0. No anxiety, at ease 1. Mildly anxious 2. 3. 4. Moderately anxious, or guarded, so anxiety inferred 5. 6. 7. Equivalent to acute panic states as seen in severe delirium or acute schizophrenic reactions 	 Some norm anorm Mode and resident Mode The second second	mal activity newhat more than mal activity0. None 1. Very mi needlesactivity1. Very mi needlesderately fidgety restless2. Mild itc needlesderately fidgety restless3. Modera needleses back and forth ng most of the rview, or stantly thrashes6. Extreme 		y mild itching, pins and dles, burning or numbness d itching, pins and edles, burning or
 7. Auditory disturbances 0. Not present 1. Very mild harshness or ability to frighten 2. Mild harshness or ability to frighten 3. Moderate harshness or ability to frighten 4. Moderately severe hallucinations 5. Severe hallucinations 6. Extremely severe hallucinations 7. Continuous hallucinations 10. Orientation — Ask: Who am I? V been happening in the commun 0. Person ✓ Place ✓ Time ✓ Orie 1. Person ✓ Place ✓ Time ✓ Disc 2. Person ✓ Place × Time × Doe 		ty? ntated, aware of community e rientated to community event		5. Severe 6. Very severe 7. Extremely severe e of day is it? Has anything nity events events vents

Observational assessment. Add up score for 10 criteria = score for person.

4. Person × Place × Time × Disorientated