



<p>This table must be used with protocols from CARPA STM (8th ed) or WBMI (7th ed) – it does not provide all the information needed for appropriate treatment † = other strengths and forms available. Doses in brackets (mL, tab) only apply to forms and strengths listed. Doses (mL) rounded up to nearest 0.2mL unless this is more than 10% above recommended dose.</p>												
Medicine and presentation	Common uses	Route and frequency	Dosage	Amount per dose							Notes	
				New-born 3.3kg	3 months 6.2kg	6 months 7.6kg	1 year 9kg	2 years 12kg	4 years 16kg	6 years 20kg		8 years 25kg
Pyrantel Tab: 125mg, 250mg Choc sq: 100mg Pregnancy: B2 – safe to use Breastfeed: safe to use	Threadworm	Oral Single dose Repeat dose in 2 weeks	10mg/kg/dose	N/A	100mg or 1 sq	120mg or 1 tab – 125mg	160mg or 1½ sq or 1 tab – 125mg	200mg or 2 sq or 1½ tab – 125mg	250mg or 2½ sq or 1 tab – 250mg	320mg or 3 sq or 2½ tab – 125mg	400mg or 4 sq or 3 tab – 125mg	Take with food. Tablet can be crushed and mixed with jam 40kg 400mg or 4 sq or 3 tab – 125mg 60kg 600mg or 6 sq or 5 tab – 125mg 80kg 800mg or 3 tab – 250mg 100kg+ 1g or 4 tab – 250mg
	Hookworm	Oral Once a day for 3 days										
Trimethoprim-sulfamethoxazole (co-trimoxazole) Susp: 8+40mg/mL Tab: 160+800mg Pregnancy: C – avoid use Breastfeed: caution	Bites Otitis media Skin infection UTI Soft tissue infection Nappy rash	Oral Twice a day (bd)	4+20mg/kg/dose	N/A	36mg (4.6mL)	48mg (6mL)	64mg (8mL)	80mg (10mL) or ½ tab	100mg (12.6mL)	128mg (16mL)	160mg (20mL) or 1 tab	Doses worked out using trimethoprim component. Addition of Folic Acid for prolonged or high dose courses.
	Melioidosis		6+30mg/kg/dose	N/A	30.4mg (3.8mL)	48mg (6mL)	64mg (8mL)	80mg (10mL) or ½ tab	100mg (12.6mL)	128mg (16mL)	160mg (20mL) or 1 tab	
	Water-related skin infections		8+40mg/kg/dose	N/A	24.8mg (3.2mL)	36mg (4.6mL)	48mg (6mL)	64mg (8mL)	100mg (12.6mL)	128mg (16mL)	160mg (20mL) or 1 tab	40-60kg 240mg (30mL or 1½ tab) 60+kg 320mg (40mL or 2 tab)